



DOSING &
ADMINISTRATION
GUIDE



TO LEARN MORE, PLEASE VISIT WWW.YONDELIS.COM

INDICATIONS

YONDELIS[®] is indicated for the treatment of patients with unresectable or metastatic liposarcoma or leiomyosarcoma who received a prior anthracycline-containing regimen.

IMPORTANT SAFETY INFORMATION

CONTRAINDICATIONS — YONDELIS[®] is contraindicated in patients with known severe hypersensitivity, including anaphylaxis, to trabectedin.

Please read Important Safety Information on the following pages and [click here](#) to read the full Prescribing Information for YONDELIS[®].

RECOMMENDED DOSING FOR YONDELIS® (trabectedin)

Recommended Dosage

- **The recommended dose** is 1.5 mg/m² administered as an intravenous infusion over 24 hours through a central venous line every 21 days (3 weeks), until disease progression or unacceptable toxicity.

Premedication

- **Administer dexamethasone** 20 mg intravenously 30 minutes prior to each dose of YONDELIS®

Dose Modifications for Adverse Reactions

Permanently discontinue YONDELIS® for:

- Persistent adverse reactions requiring a delay in dosing of more than 3 weeks
- Adverse reactions requiring dose reduction following YONDELIS® administered at 1.0 mg/m² for patients with normal hepatic function or at 0.3 mg/m² for patients with pre-existing moderate hepatic impairment
- Severe liver dysfunction: bilirubin 2 times the upper limit of normal and AST or ALT 3 times the upper limit of normal with ALP less than 2 times the upper limit of normal in the prior treatment cycle for patients with normal liver function at baseline
- Exacerbation of liver dysfunction in patients with pre-existing moderate hepatic impairment
- Capillary leak syndrome
- Rhabdomyolysis
- Grade 3 or 4 cardiac adverse events (AEs) indicative of cardiomyopathy or for subjects with an LVEF that decreases below the lower limit of normal

The recommended dose modifications for specific adverse reactions are listed in the table on the following page. Once reduced, the dose of YONDELIS® should not be increased in subsequent treatment cycles.

Hepatic Impairment: The recommended dose is 0.9 mg/m² in patients with moderate hepatic impairment (bilirubin levels greater than 1.5 times to 3 times the upper limit of normal, and AST and ALT less than 8 times the upper limit of normal). Do not administer YONDELIS® to patients with severe hepatic impairment (bilirubin levels above 3 times the upper limit of normal, and any AST and ALT) [see *Use in Specific Populations (8.6), Clinical Pharmacology (12.3)*].

Please read Important Safety Information on the following pages and [click here](#) to read the full Prescribing Information for YONDELIS®.

Recommended Starting Doses and Dose Reductions

Starting Dose and Dose Reduction	For patients with normal hepatic function or mild hepatic impairment* prior to initiation of YONDELIS® treatment	For patients with moderate hepatic impairment† prior to initiation of YONDELIS® treatment
Starting dose	1.5 mg/m ²	0.9 mg/m ²
Dose reduction		
First dose reduction	1.2 mg/m ²	0.6 mg/m ²
Second dose reduction	1.0 mg/m ²	0.3 mg/m ²

*Including patients with bilirubin greater than 1 to 1.5 times the upper limit of normal and any AST or ALT.

†Including patients with bilirubin levels greater than 1.5 times to 3 times the upper limit of normal and AST and ALT less than 8 times the upper limit of normal.

Recommended Dose Modifications

Laboratory Result or Adverse Reaction	DELAY Next Dose for Up to 3 Weeks	REDUCE Next Dose by 1 Dose Level for Adverse Reaction(s) During Prior Cycle
Platelets	<100,000 platelets/microliter	<25,000 platelets/microliter
Absolute neutrophil count	<1,500 neutrophils/microliter	<1,000 neutrophils/microliter with fever/infection <500 cells/microliter lasting more than 5 days
Total bilirubin	>upper limit of normal	>upper limit of normal
AST or ALT	>2.5 times the upper limit of normal	>5 times the upper limit of normal
ALP	>2.5 times the upper limit of normal	>2.5 times the upper limit of normal
CPK	>2.5 times the upper limit of normal	>5 times the upper limit of normal
Other non-hematologic adverse reactions	Grade 3 or 4	Grade 3 or 4

ALP = alkaline phosphatase; ALT = alanine aminotransferase;
AST = aspartate aminotransferase; CPK = creatine phosphokinase.

Please read Important Safety Information on the following pages and [click here](#) to read the full Prescribing Information for YONDELIS®.



IMPORTANT SAFETY INFORMATION (cont'd)

WARNINGS AND PRECAUTIONS

Neutropenic sepsis, including fatal cases, can occur. In Trial ET743-SAR-3007, the incidence of Grade 3 or 4 neutropenia, based on laboratory values, was 43% (161/378). Median time to the first occurrence of Grade 3 or 4 neutropenia was 16 days (range: 8 days to 9.7 months). Median time to complete resolution of neutropenia was 13 days (range: 3 days to 2.3 months). Febrile neutropenia (fever $\geq 38.5^{\circ}\text{C}$ with Grade 3 or 4 neutropenia) occurred in 18 patients (5%) treated with YONDELIS[®]. Ten patients (2.6%) experienced neutropenic sepsis, 5 of whom had febrile neutropenia, which was fatal in 4 patients (1.1%). Assess neutrophil count prior to administration of each dose of YONDELIS[®] and periodically throughout the treatment cycle. Withhold or reduce dose of YONDELIS[®] based on severity of adverse reaction.

Rhabdomyolysis — YONDELIS[®] can cause rhabdomyolysis and musculoskeletal toxicity. In Trial ET743-SAR-3007, rhabdomyolysis leading to death occurred in 3 (0.8%) of the 378 patients receiving YONDELIS[®]. Elevations in creatine phosphokinase (CPK) occurred in 122 (32%) of the 378 patients receiving YONDELIS[®], including Grade 3 or 4 CPK elevation in 24 patients (6%), compared to 15 (9%) of the 172 patients receiving dacarbazine with any CPK elevation, including 1 patient (0.6%) with Grade 3 CPK elevation. Among the 24 patients receiving YONDELIS[®] with Grade 3 or 4 CPK elevation, renal failure occurred in 11 patients (2.9%); rhabdomyolysis with the complication of renal failure occurred in 4 of these 11 patients (1.1%). Median time to first occurrence of Grade 3 or 4 CPK elevations was 2 months (range: 1 to 11.5 months). Median time to complete resolution was 14 days (range: 5 days to 1 month). Assess CPK levels prior to each administration of YONDELIS[®]. Withhold, reduce dose, or permanently discontinue based on severity of adverse reaction.

Hepatotoxicity, including hepatic failure, can occur. Patients with serum bilirubin levels above the upper limit of normal or AST or ALT levels >2.5 x upper limit of normal were not enrolled in Trial ET743-SAR-3007. In Trial ET743-SAR-3007, the incidence of Grade 3-4 elevated liver function tests (defined as elevations in ALT, AST, total bilirubin, or alkaline phosphatase) was 35% (134/378) in patients receiving YONDELIS[®]. Median time to development of Grade 3-4 elevation in ALT or AST was 29 days (range: 3 days to 11.5 months). Of the 134 patients with Grade 3 to 4 elevations in LFTs, 114 (85%) experienced complete resolution with the median time to complete resolution of 13 days (range: 4 days to 4.4 months). In Trial ET743-SAR-3007, the incidence of drug-induced liver injury (defined as concurrent elevation in ALT or AST of more than three times the upper limit of normal, alkaline phosphatase less than two times the upper limit of normal, and total bilirubin at least two times the upper limit of normal) was 1.3% (5/378) in patients receiving YONDELIS[®]. ALT or AST elevation greater than eight times the upper limit of

normal occurred in 18% (67/378) of patients receiving YONDELIS®. Assess LFTs prior to each administration of YONDELIS® and as clinically indicated based on underlying severity of pre-existing hepatic impairment. Manage elevated LFTs with treatment interruption, dose reduction, or permanent discontinuation based on severity and duration of LFT abnormality.

PREPARING AND ADMINISTERING YONDELIS®

YONDELIS® is a cytotoxic drug. Follow applicable special handling and disposal procedures.

Reconstitute

- Using aseptic technique, inject 20 mL of Sterile Water for Injection, USP into the vial. Shake the vial until complete dissolution. The reconstituted solution is clear, colorless to pale brownish-yellow, and contains 0.05 mg/mL of trabectedin
- Inspect for particulate matter and discoloration prior to further dilution. Discard vial if particles or discoloration are observed

Dilute

- Immediately following reconstitution, withdraw the calculated volume of trabectedin and further dilute in 500 mL of 0.9% Sodium Chloride, USP or 5% Dextrose Injection, USP
- Do not mix YONDELIS® with other drugs

Administer

- Infuse the reconstituted, diluted solution over 24 hours through a central venous line using an infusion set with a 0.2-micron polyethersulfone (PES) in-line filter to reduce the risk of exposure to adventitious pathogens that may be introduced during solution preparation
- Complete infusion within 30 hours of initial reconstitution

Discard

- Discard any remaining solution within 30 hours of reconstituting the lyophilized powder

Please read Important Safety Information on the following pages and [click here](#) to read the full Prescribing Information for YONDELIS®.



Compatibility

- YONDELIS® diluted solution is compatible with Type I colorless glass vials, polyvinylchloride (PVC) and polyethylene (PE) bags and tubing, PE and polypropylene (PP) mixture bags, polyethersulfone (PES) in-line filters, titanium, platinum or plastic ports, silicone and polyurethane catheters, and pumps having contact surfaces made of PVC, PE, or PE/PP

IMPORTANT SAFETY INFORMATION (cont'd)

WARNINGS AND PRECAUTIONS (cont'd)

Cardiomyopathy, including cardiac failure, congestive heart failure, ejection fraction decreased, diastolic dysfunction, or right ventricular dysfunction can occur. In Trial ET743-SAR-3007, a significant decrease in left ventricular ejection fraction (LVEF) was defined as an absolute decrease of $\geq 15\%$ or below the lower limit of normal with an absolute decrease of $\geq 5\%$. Patients with a history of New York Heart Association Class II to IV heart failure or abnormal LVEF at baseline were ineligible. In Trial ET743-SAR-3007, cardiomyopathy occurred in 23 patients (6%) receiving YONDELIS® and in four patients (2.3%) receiving dacarbazine. Grade 3 or 4 cardiomyopathy occurred in 15 patients (4%) receiving YONDELIS® and 2 patients (1.2%) receiving dacarbazine; cardiomyopathy leading to death occurred in 1 patient (0.3%) receiving YONDELIS® and in none of the patients receiving dacarbazine. The median time to development of Grade 3 or 4 cardiomyopathy in patients receiving YONDELIS® was 5.3 months (range: 26 days to 15.3 months). Patients with LVEF < lower limit of normal, prior cumulative anthracycline dose of ≥ 300 mg/m², age ≥ 65 years, or a history of cardiovascular disease may be at increased risk of cardiac dysfunction. Assess LVEF by echocardiogram (ECHO) or multigated acquisition (MUGA) scan before initiation of YONDELIS® and at 2- to 3-month intervals thereafter until YONDELIS® is discontinued. Discontinue treatment with YONDELIS® based on severity of adverse reaction.

For additional information, please read the full Prescribing Information.

Please read Important Safety Information on the following pages and [click here](#) to read the full Prescribing Information for YONDELIS®.

HANDLING YONDELIS®

How supplied

- YONDELIS® is supplied in a single-dose glass vial containing 1 mg trabectedin
- Each outer carton of YONDELIS® contains 1 vial
- NDC: 59676-610-01

Storage and handling

Store in a refrigerator at 2°C to 8°C (36°F to 46°F).

YONDELIS® is a cytotoxic drug. Follow applicable special handling and disposal procedures.



Bottle shown is not actual size.

IMPORTANT SAFETY INFORMATION

CONTRAINDICATIONS — YONDELIS® (trabectedin) is contraindicated in patients with known severe hypersensitivity, including anaphylaxis, to trabectedin.

WARNINGS AND PRECAUTIONS

Neutropenic sepsis, including fatal cases, can occur. In Trial ET743-SAR-3007, the incidence of Grade 3 or 4 neutropenia, based on laboratory values, was 43% (161/378). Median time to the first occurrence of Grade 3 or 4 neutropenia was 16 days (range: 8 days to 9.7 months). Median time to complete resolution of neutropenia was 13 days (range: 3 days to 2.3 months). Febrile neutropenia (fever $\geq 38.5^{\circ}\text{C}$ with Grade 3 or 4 neutropenia) occurred in 18 patients (5%) treated with YONDELIS®. Ten patients (2.6%) experienced neutropenic sepsis, 5 of whom had febrile neutropenia, which was fatal in 4 patients (1.1%). Assess neutrophil count prior to administration of each dose of YONDELIS® and periodically throughout the treatment cycle. Withhold or reduce dose of YONDELIS® based on severity of adverse reaction.

Rhabdomyolysis — YONDELIS® can cause rhabdomyolysis and musculoskeletal toxicity. In Trial ET743-SAR-3007, rhabdomyolysis leading to death occurred in 3 (0.8%) of the 378 patients receiving YONDELIS®. Elevations in creatine phosphokinase (CPK) occurred in 122 (32%) of the 378 patients receiving YONDELIS®, including Grade 3 or 4 CPK elevation in 24 patients (6%), compared to 15 (9%) of the 172 patients receiving dacarbazine with any CPK elevation, including 1 patient (0.6%) with Grade 3 CPK elevation. Among the 24 patients receiving YONDELIS® with Grade 3 or 4 CPK elevation, renal failure occurred in 11 patients (2.9%); rhabdomyolysis with the complication of renal failure occurred in 4 of these 11 patients (1.1%). Median time to first occurrence of Grade 3 or 4 CPK elevations was 2 months (range: 1 to 11.5 months). Median time to complete resolution was 14 days (range: 5 days to 1 month). Assess CPK levels prior to each administration of YONDELIS®. Withhold, reduce dose, or permanently discontinue based on severity of adverse reaction.

IMPORTANT SAFETY INFORMATION (cont'd)

Hepatotoxicity, including hepatic failure, can occur. Patients with serum bilirubin levels above the upper limit of normal or AST or ALT levels >2.5 x upper limit of normal were not enrolled in Trial ET743-SAR-3007. In Trial ET743-SAR-3007, the incidence of Grade 3-4 elevated liver function tests (defined as elevations in ALT, AST, total bilirubin, or alkaline phosphatase) was 35% (134/378) in patients receiving YONDELIS[®]. Median time to development of Grade 3-4 elevation in ALT or AST was 29 days (range: 3 days to 11.5 months). Of the 134 patients with Grade 3 to 4 elevations in LFTs, 114 (85%) experienced complete resolution with the median time to complete resolution of 13 days (range: 4 days to 4.4 months). In Trial ET743-SAR-3007, the incidence of drug-induced liver injury (defined as concurrent elevation in ALT or AST of more than three times the upper limit of normal, alkaline phosphatase less than two times the upper limit of normal, and total bilirubin at least two times the upper limit of normal) was 1.3% (5/378) in patients receiving YONDELIS[®]. ALT or AST elevation greater than eight times the upper limit of normal occurred in 18% (67/378) of patients receiving YONDELIS[®]. Assess LFTs prior to each administration of YONDELIS[®] and as clinically indicated based on underlying severity of pre-existing hepatic impairment. Manage elevated LFTs with treatment interruption, dose reduction, or permanent discontinuation based on severity and duration of LFT abnormality.

Cardiomyopathy, including cardiac failure, congestive heart failure, ejection fraction decreased, diastolic dysfunction, or right ventricular dysfunction can occur. In Trial ET743-SAR-3007, a significant decrease in left ventricular ejection fraction (LVEF) was defined as an absolute decrease of $\geq 15\%$ or below the lower limit of normal with an absolute decrease of $\geq 5\%$. Patients with a history of New York Heart Association Class II to IV heart failure or abnormal LVEF at baseline were ineligible. In Trial ET743-SAR-3007, cardiomyopathy occurred in 23 patients (6%) receiving YONDELIS[®] and in four patients (2.3%) receiving dacarbazine. Grade 3 or 4 cardiomyopathy occurred in 15 patients (4%) receiving YONDELIS[®] and 2 patients (1.2%) receiving dacarbazine; cardiomyopathy leading to death occurred in 1 patient (0.3%) receiving YONDELIS[®] and in none of the patients receiving dacarbazine. The median time to development of Grade 3 or 4 cardiomyopathy in patients receiving YONDELIS[®] was 5.3 months (range: 26 days to 15.3 months). Patients with LVEF $<$ lower limit of normal, prior cumulative anthracycline dose of ≥ 300 mg/m², age ≥ 65 years, or a history of cardiovascular disease may be at increased risk of cardiac

[Click here to read the full Prescribing Information.](#)



IMPORTANT SAFETY INFORMATION (cont'd)

dysfunction. Assess LVEF by echocardiogram (ECHO) or multigated acquisition (MUGA) scan before initiation of YONDELIS® and at 2- to 3-month intervals thereafter until YONDELIS® is discontinued. Discontinue treatment with YONDELIS® based on severity of adverse reaction.

Capillary leak syndrome (CLS) characterized by hypotension, edema, and hypoalbuminemia has been reported with YONDELIS®, including serious CLS resulting in death. Monitor for signs and symptoms of CLS. Discontinue YONDELIS® and promptly initiate standard management for patients with CLS, which may include a need for intensive care.

Extravasation Resulting in Tissue Necrosis — Extravasation of YONDELIS®, resulting in tissue necrosis requiring debridement, can occur. Evidence of tissue necrosis can occur more than 1 week after the extravasation. There is no specific antidote for extravasation of YONDELIS®. Administer YONDELIS® through a central venous line.

Embryo-Fetal Toxicity — Based on its mechanism of action, YONDELIS® can cause fetal harm when administered to a pregnant woman. Advise females of reproductive potential to use effective contraception during therapy and for at least 2 months after the last dose of YONDELIS®. Advise males with female partners of reproductive potential to use effective contraception during therapy and for at least 5 months after the last dose of YONDELIS®.

Adverse Reactions — The most common ($\geq 20\%$) adverse reactions are nausea (75%), fatigue (69%), vomiting (46%), constipation (37%), decreased appetite (37%), diarrhea (35%), peripheral edema (28%), dyspnea (25%), and headache (25%).

The most common ($\geq 5\%$) grades 3-4 laboratory abnormalities are: neutropenia (43%), increased ALT (31%), thrombocytopenia (21%), anemia (19%), increased AST (17%), and increased creatine phosphokinase (6.4%).

IMPORTANT SAFETY INFORMATION (cont'd)

DRUG INTERACTIONS

Effect of Cytochrome CYP3A Inhibitors — Avoid using strong CYP3A inhibitors (e.g., oral ketoconazole, itraconazole, posaconazole, voriconazole, clarithromycin, telithromycin, indinavir, lopinavir, ritonavir, boceprevir, nelfinavir, saquinavir, telaprevir, nefazodone, conivaptan) in patients taking YONDELIS®. If a strong CYP3A inhibitor for short-term use (i.e., less than 14 days) must be used, administer the strong CYP3A inhibitor 1 week after the YONDELIS® infusion, and discontinue it the day prior to the next YONDELIS® infusion.

Effect of Cytochrome CYP3A Inducers — Avoid using strong CYP3A inducers (e.g., rifampin, phenobarbital, St. John's wort) in patients taking YONDELIS®.

[Click here](#) to read the full Prescribing Information for YONDELIS®.

cp-63569v2

Safety





Have any questions about
YONDELIS® (trabectedin)?



Call Janssen Medical Information Center
1-800-JANSSEN (1-800-526-7736)



Or visit **janssenmd.com**

TO LEARN MORE, PLEASE VISIT **WWW.YONDELIS.COM**



Please read Important Safety Information on the following pages and [click here](#) to read the full Prescribing Information for YONDELIS®.

